



A LETTERS JOURNAL EXPLORING
THE FRONTIERS OF PHYSICS

OFFPRINT

Thermal conductance of the coupled-rotator chain: Influence of temperature and size

YUNYUN LI, NIANBEI LI, UGUR TIRNAKLI BAOWEN LI and
CONSTANTINO TSALLIS

EPL, **117** (2017) 60004

Please visit the website
www.epljournal.org

Note that the author(s) has the following rights:

- immediately after publication, to use all or part of the article without revision or modification, **including the EPLA-formatted version**, for personal compilations and use only;
- no sooner than 12 months from the date of first publication, to include the accepted manuscript (all or part), **but not the EPLA-formatted version**, on institute repositories or third-party websites provided a link to the online EPL abstract or EPL homepage is included.

For complete copyright details see: <https://authors.epljournal.net/documents/copyright.pdf>.



epl

A LETTERS JOURNAL EXPLORING
THE FRONTIERS OF PHYSICS

AN INVITATION TO SUBMIT YOUR WORK

epljournal.org

The Editorial Board invites you to submit your letters to EPL

EPL is a leading international journal publishing original, innovative Letters in all areas of physics, ranging from condensed matter topics and interdisciplinary research to astrophysics, geophysics, plasma and fusion sciences, including those with application potential.

The high profile of the journal combined with the excellent scientific quality of the articles ensures that EPL is an essential resource for its worldwide audience. EPL offers authors global visibility and a great opportunity to share their work with others across the whole of the physics community.

Run by active scientists, for scientists

EPL is reviewed by scientists for scientists, to serve and support the international scientific community. The Editorial Board is a team of active research scientists with an expert understanding of the needs of both authors and researchers.



epljournal.org

OVER

568,000

full text downloads in 2015

18 DAYS

average accept to online
publication in 2015

20,300

citations in 2015

*"We greatly appreciate
the efficient, professional
and rapid processing of
our paper by your team."*

Cong Lin
Shanghai University

Six good reasons to publish with EPL

We want to work with you to gain recognition for your research through worldwide visibility and high citations. As an EPL author, you will benefit from:

- 1 Quality** – The 60+ Co-editors, who are experts in their field, oversee the entire peer-review process, from selection of the referees to making all final acceptance decisions.
- 2 Convenience** – Easy to access compilations of recent articles in specific narrow fields available on the website.
- 3 Speed of processing** – We aim to provide you with a quick and efficient service; the median time from submission to online publication is under 100 days.
- 4 High visibility** – Strong promotion and visibility through material available at over 300 events annually, distributed via e-mail, and targeted mailshot newsletters.
- 5 International reach** – Over 3200 institutions have access to EPL, enabling your work to be read by your peers in 100 countries.
- 6 Open access** – Articles are offered open access for a one-off author payment; green open access on all others with a 12-month embargo.

Details on preparing, submitting and tracking the progress of your manuscript from submission to acceptance are available on the EPL submission website epletters.net.

If you would like further information about our author service or EPL in general, please visit epijournal.org or e-mail us at info@epijournal.org.

EPL is published in partnership with:



European Physical Society



Società Italiana
di Fisica

Società Italiana di Fisica

 **edp sciences** **IOP Publishing**

EDP Sciences

IOP Publishing

Thermal conductance of the coupled-rotator chain: Influence of temperature and size

YUNYUN LI^{1,2,3}, NIANBEI LI^{1,2,3}, UGUR TIRNAKLI^{4,5} BAOWEN LI⁶ and CONSTANTINO TSALLIS^{5,7,8}

¹Center for Phononics and Thermal Energy Science and School of Physics Science and Engineering, Tongji University 200092 Shanghai, China

²China-EU Joint Lab for Nanophononics, Tongji University - Shanghai 200092, China

³Shanghai Key Laboratory of Special Artificial Microstructure Materials and Technology, School of Physics Science and Engineering, Tongji University - Shanghai 200092, China

⁴Department of Physics, Faculty of Science, Ege University - 35100 Izmir, Turkey

⁵Centro Brasileiro de Pesquisas Físicas and National Institute for Science and Technology of Complex Systems Rua Dr. Xavier Sigaud 150, 22290-180 Rio de Janeiro, RJ, Brazil

⁶Department of Mechanical Engineering, University of Colorado Boulder - Boulder, CO 80309, USA

⁷Santa Fe Institute - 1399 Hyde Park Road, Santa Fe, NM 87501, USA

⁸Complexity Science Hub Vienna - Josefstadter Strasse 39, 1080 Vienna, Austria

received 22 March 2017; accepted in final form 3 May 2017

published online 18 May 2017

PACS 05.20.-y – Classical statistical mechanics

PACS 05.45.Pq – Numerical simulations of chaotic systems

Abstract – The thermal conductance of a homogeneous 1D nonlinear lattice system with nearest-neighbor interactions has recently been computationally studied in detail by Li *et al.* (*Eur. Phys. J. B*, **88** (2015) 182), where its power-law dependence on temperature T for high temperatures is shown. Here, we address its entire temperature dependence, in addition to its dependence on the size N of the system. We obtain a neat data collapse for arbitrary temperatures and system sizes, and numerically show that the thermal conductance curve is quite satisfactorily described by a fat-tailed q -Gaussian dependence on $TN^{1/3}$ with $q \simeq 1.55$. Consequently, its $T \rightarrow \infty$ asymptotic behavior is given by $T^{-\alpha}$ with $\alpha = 2/(q-1) \simeq 3.64$.

Copyright © EPLA, 2017

Introduction. – The breakdown of Fourier's law in low-dimensional lattices has attracted much attention in recent years due to its fundamental importance within non-equilibrium thermodynamics and statistical mechanics [1–6]. In the 1D Fermi-Pasta-Ulam (FPU- β) lattice, the thermal conductivities κ diverge with system sizes N as $\kappa \propto N^\gamma$, where $0 < \gamma < 1$; consequently its thermal conductance $\sigma \equiv \kappa/N$ vanishes in the $N \rightarrow \infty$ limit. However, there is still no clear conclusion about the physical ingredient responsible for this kind of anomalous heat conduction. It is believed that momentum conservation is the crucial reason for the anomalous heat conduction [7–9], but normal heat conduction has been found in the 1D coupled rotator lattice, which also is a momentum-conserved system [10,11].

Unlike the FPU- β -like lattices, the 1D coupled rotator lattice has periodic interatomic potential which is finite. As a result, the energy diffusion as well as the momentum

diffusion are normal [12]. In order to understand the effect of this finite interatomic potential, previous works focus on the temperature dependence of the thermal conductivities in the 1D coupled rotator lattice [10,11]. In both works, the thermal conductivity was proposed to have an exponential dependence on temperature. But it is argued that $\kappa(T) \propto e^{\Delta V/T}$, where ΔV is proportional to the potential barrier height in [10], while $\kappa(T) \propto e^{-T/A}$ with A a fitting parameter in [11]. It has only recently been found that the temperature dependence in the 1D coupled rotator lattice follows a power-law behavior on temperature as $\kappa(T) \propto T^{-\alpha}$ with $\alpha \simeq 3.2$ for intermediate temperatures [13]. Interestingly enough, this power-law dependence is qualitatively consistent with the theoretical prediction for the Chirikov standard map which is a single-rotator model [14,15].

On the other hand, the standard map, as well as several other dynamical complex systems, has recently

been shown [16] to present non-Gaussian probability distributions for the sum of its position random variable. These distributions are approached extremely well by the q -Gaussian defined as

$$P_q(x) = A_q e_q^{-B_q x^2} \equiv \frac{A_q}{[1 + (q-1)B_q x^2]^{1/(q-1)}}, \quad (1)$$

where A_q is the normalization factor, $B_q > 0$ is a parameter which characterizes the width of the distribution, and the index $q \geq 1$ [17–19]. In the $q \rightarrow 1$ limit, this expression recovers the standard Gaussian distribution. This family of distributions optimizes the nonadditive entropy $S_q = k \frac{1 - \int dx [p(x)]^q}{q-1}$ with $S_1 = S_{BG} \equiv -k \int dx p(x) \ln p(x)$ under appropriate constraints, where k is the Boltzmann constant, $p(x)$ is the probability distribution, and BG stands for Boltzmann-Gibbs.

Model and method. – In this letter, we study a homogeneous 1D nonlinear lattice system with nearest-neighbor interactions and try to see whether some of its properties also are consistent with q -Gaussians. For this lattice system, the Hamiltonian with the corresponding dimensionless units can be written in the general form

$$H = \sum_{i=1}^N \left[\frac{p_i^2}{2} + V(q_{i+1} - q_i) + U(q_i) \right], \quad (2)$$

where p_i denotes the momentum for the i -th rotator. The set q_i are the displacements from the equilibrium position for the i -th rotator; $V(q_{i+1} - q_i)$ is the interaction potential between neighboring sites i and $i+1$, and $U(q_i)$ is the on-site potential, representing the interaction with the substrate. To focus on the momentum-conserving system, we set $U = 0$. The potential we employ is in the form

$$V(x) = V_0(1 - \cos x), \quad (3)$$

where V_0 is the interaction strength (without loss of generality we set $V_0 = 1$).

In our simulation, a Langevin form of heat bath is used. For the chain with N particles, only the first and last particles are coupled to the heat bath, with the temperature T_L and T_R , respectively. The dynamics equations of the motion read

$$\begin{aligned} \dot{q}_i &= p_i, \quad i = 1, 2, 3, \dots, N, \\ \dot{p}_i &= F(q_i - q_{i-1}) + F(q_{i+1} - q_i), \quad i = 2, 3, \dots, N-1, \\ \dot{p}_1 &= F(q_1) + F(q_{i+1} - q_i) - \gamma p_1 + \xi_1, \\ \dot{p}_N &= F(q_i - q_{i-1}) + F(-q_i) - \gamma p_N + \xi_N, \end{aligned} \quad (4)$$

where $F(x) = -\partial V(x)/\partial q_i$ and γ is the friction coefficient; ξ_1, ξ_N stand for the Gaussian white noise with zero mean $\langle \xi_1(t) \rangle = 0$ and $\langle \xi_N(t) \rangle = 0$. The correlation functions are given by

$$\begin{aligned} \langle \xi_1(t) \xi_1(t') \rangle &= 2\gamma T_L \delta(t - t'), \\ \langle \xi_N(t) \xi_N(t') \rangle &= 2\gamma T_R \delta(t - t'). \end{aligned} \quad (5)$$




Fig. 1: (Color online) Heat flow at stationary state for typical lattice sizes. Top: thermal conductivity $\kappa \equiv \sigma N$ (notice that data collapse occurs for the high-temperature region). Bottom: thermal conductance σ (notice that data collapse occurs for the low-temperature region). Different colors correspond to the lattice lengths $N = 50, 100, 200, 400, 800$ and 2000 . The slope -3.2 indicated in [13] is shown here for comparison. The dashed curves are guides to the eye.

For simplicity, the temperatures are set as $T_{L/R} = T_0(1 \pm \Delta)$, where T_0 is the average temperature and Δ is the temperature difference. Throughout our numerical simulations, $\Delta = 0.1$ is restricted to the small perturbation regime and $\gamma = 1$ is fixed. The evolution of dynamics (eqs. (4)) is integrated by the Verlet velocity algorithm and the time step $\Delta t = 0.01$, which is small enough [20]. All the results are analyzed for the time scale 10^7 – 10^8 , after the system release to the steady state.

Results and discussion. – The thermal conductivity κ is characterized by

$$\kappa(T) = \frac{JN}{T_L - T_R}, \quad (6)$$

where $J = \langle J_i \rangle$ is the average heat flux along the lattice and J_i is the local heat flux. As already mentioned, the thermal conductance is defined as $\sigma \equiv \kappa/N$. The temperature dependence of the thermal conductance is given in fig. 1 for six different lattice sizes. The asymptotic power-law behavior is evident with an exponent -3.2 . One can easily obtain a clear data collapse as shown in fig. 2. It is evident that the temperature dependence of the




Fig. 2: (Color online) Data collapse for lattice sizes going from $N = 25$ to $N = 2000$. The continuous curve (green line) corresponds to $\sigma = A_q e_q^{-B_q (TN^{1/3})^2}$ with $(q, B_q, A_q) = (1.55, 0.40, 0.189)$. Notice that the maximum value 0.189 has naturally nothing to do with the normalizing factor of a q -Gaussian distribution, which for the present values $(q, B_q) = (1.55, 0.40)$ would approximately be 0.28. Indeed, q -exponential and q -Gaussian functions frequently appear in diverse physical quantities (*e.g.*, the time evolution of the non-linear dynamical sensitivity to the initial conditions [21,22]), and not only as probability distributions. The asymptotic slope is given by $2/(1-q) \simeq -3.64$, in contrast with the intermediate slope -3.2 indicated in [13] (see fig. 1). Notice that the asymptotic slope only becomes visible several decades further down.

thermal conductance can be satisfactorily approached by a q -Gaussian with $q = 1.55$. At this point, it is worth mentioning that another system which can be considered to be similar to ours has already been shown to exhibit strong deviations from the Gaussian behavior [23]. In that contribution, the authors analytically study a Brownian particle with linear-position-dependent inverse temperature with a confining potential and show that the behavior can be well described by superstatistics [24], which includes q -Gaussians as a special case.

In conclusion, we have numerically determined that the thermal conductance for the classical one-dimensional first-neighbor coupled planar-rotator (or *inertial XY ferromagnetic*) chain (eqs. (2) and (3), with vanishing on-site potential $U(q_i)$ and unit potential strength V_0) is amazingly well described by the q -Gaussian $\sigma \propto e_{1.55}^{-0.40(TN^{1/3})^2}$ for wide ranges of temperature T and lattice size N . This result implies that, in the $(TN^{1/3}) \rightarrow \infty$ limit, we asymptotically expect $\sigma \propto [TN^{1/3}]^{-\alpha}$ with $\alpha = 2/(q-1) \simeq 3.64$, close yet different from the value 3.2 determined in [13] for intermediate temperatures. At thermal equilibrium (*i.e.*, for $T_L = T_R$), it is clear that the present short-range interacting model follows the Boltzmann-Gibbs statistical mechanics. Why then, in the nonequilibrium stationary state characterized by $T_L \neq T_R$, does such a strong suggestion of q -statistics emerge? This remains as a highly interesting and certainly intriguing open question, somewhat reminiscent of the aging and related phenomena in

various systems: see for example [25] (its fig. 1), [26], and [27] (its fig. 1, for instance); see also [28] (its fig. 3). It is of course not excluded that, due to the permanent unidirectional heat flow, the phase space of the chain is visited in an incomplete manner. Such type of behavior could be related to the frequent emergence of q -statistics in hydrology [29]. As a final remark let us consider at this point why the distribution that we obtain is a q -Gaussian and not a q -exponential. This is possibly due to the fact that the heat flow occurs always from the hot source to the cold one. This fact results in a symmetry which is very reminiscent of the distribution observed in [30] where the rotation periods of asteroids are found to follow q -Gaussians, in contrast with asteroid sizes, observed to follow a q -exponential. Indeed, the periods are related to the angular velocities, which can rotate in both senses. Further understanding would of course be very welcome.

YL, NL and BL are supported by the NSF China with grant No. 11334007. UT is a member of the Science Academy, Istanbul, Turkey. Two of us (UT and CT) acknowledge partial financial support from CNPq, Capes and Faperj (Brazilian agencies), as well as from the John Templeton Foundation (USA).

REFERENCES

- [1] LEPRI S., LIVI R. and POLITI A., *Phys. Rep.*, **377** (2003) 1.
- [2] DHAR A., *Adv. Phys.*, **57** (2008) 457.
- [3] LIU S., XU X., XIE R., ZHANG G. and LI B., *Eur. Phys. J. B*, **85** (2012) 337.
- [4] CHEN S. D., WANG J., CASATI G. and BENENTI G., *Phys. Rev. E*, **90** (2014) 032134.
- [5] OLIVARES C. and ANTENEODO C., *Phys. Rev. E*, **94** (2016) 042117.
- [6] BAGCHI D., *Phys. Rev. E*, **95** (2017) 032102.
- [7] HU B., LI B. and ZHAO H., *Phys. Rev. E*, **57** (1998) 2992.
- [8] PROSEN T. and CAMPBELL D. K., *Phys. Rev. Lett.*, **84** (2000) 2857.
- [9] NARAYAN O. and RAMASWAMY S., *Phys. Rev. Lett.*, **89** (2002) 200601.
- [10] GIARDINÁ C., LIVI R., POLITI A. and VASSALLI M., *Phys. Rev. Lett.*, **84** (2000) 2144.
- [11] GENDELMAN O. and SAVIN A., *Phys. Rev. Lett.*, **84** (2000) 2381.
- [12] LI Y., LIU S., LI N., HÄNGGI P. and LI B., *New J. Phys.*, **17** (2015) 043064.
- [13] LI Y., LI N. and LI B., *Eur. Phys. J. B*, **88** (2015) 182.
- [14] CHIRIKOV B. V., *Phys. Rep.*, **52** (1979) 263.
- [15] MACKAY R. S., MEISS J. D. and PERCIVAL I. C., *Physica D*, **13** (1984) 55.
- [16] TIRNAKLI U. and BORGES E. P., *Sci. Rep.*, **6** (2016) 23644.
- [17] TSALLIS C., *J. Stat. Phys.*, **52** (1988) 479.
- [18] PRATO D. and TSALLIS C., *Phys. Rev. E*, **60** (1999) 2398.

- [19] TSALLIS C., *Introduction to Nonextensive Statistical Mechanics - Approaching a Complex World* (Springer, New York) 2009.
- [20] VERLET L., *Phys. Rev.*, **159** (1967) 98.
- [21] COSTA U. M. S., LYRA M. L., PLASTINO A. R. and TSALLIS C., *Phys. Rev. E*, **56** (1997) 245.
- [22] LYRA M. L. and TSALLIS C., *Phys. Rev. Lett.*, **80** (1998) 53.
- [23] VAN DER STRAETEN E. and BECK C., *Chin. Sci. Bull.*, **56** (2011) 3633.
- [24] BECK C. and COHEN E. G. D., *Physica A*, **322** (2003) 267.
- [25] CUGLIANDOLO L. F. and DEAN D. S., *J. Phys. A*, **28** (1995) 4213.
- [26] STARIOLO D. A., MONTEMURRO M. A. and TAMARIT F. A., *Eur. Phys. J. B*, **32** (2003) 361.
- [27] FYODOROV Y. V., PERRET A. and SCHEHR G., *J. Stat. Mech.* (2015) P11017.
- [28] DOYE J. P. K. and MASSEN C. P., *J. Chem. Phys.*, **122** (2005) 084105.
- [29] SINGH V. P., *Introduction to Tsallis Entropy Theory in Water Engineering* (CRC Press, Taylor and Francis) 2016.
- [30] BETZLER A. S. and BORGES E. P., *Astron. Astrophys.*, **539** (2012) A158.